

Optical characterization of InAs/InP quantum dots optically-active in the second telecom window

Anna Penkala¹, Paweł Podemski¹, Vinayakrishna Joshi², Vitalii Sichkovskiy², Johann P. Reithmaier² and Grzegorz Sęk¹

¹ *Department of Experimental Physics, Faculty of Fundamental Problems of Technology, Wrocław University of Science and Technology, Wybrzeże Stanisława Wyspiańskiego 27, 50-370 Wrocław, Poland*

² *Technological Physics, Institute of Nanostructure Technologies and Analytics, CINSaT, University of Kassel, 34132 Kassel, Germany*

For the realization of optical fiber communication, the O-band (original band: 1260-1360 nm) provides low transmission losses, offering at the same time very weak chromatic dispersion, limiting undesirable effects of pulse broadening and chirping. Here, we look at the InAs quantum dots developed as the active region of O-band-compatible lasers. The structures were grown by molecular-beam epitaxy on InP substrate, which became an universal material platform for quantum dots optically-active in O-band and C-band. One of the expected active region features are limited carrier losses, so the studied structures were grown with different compositions of InAlGaAs barrier surrounding active region, which was also grown with different number of quantum dot layers.




Fig.1. Example of photoreflectance signal analysis in the spectral range of quantum dot transitions

To derive the information on the energetic structure we use a few optical spectroscopy techniques: photoluminescence, photoreflectance and photoluminescence excitation. Photoluminescence spectra provide general emission properties and, in some cases, the excited states can also be observed. Photoreflectance is based on reflectance measurement where the electric field within the structure is periodically modulated by a laser beam. It gives information on the overall energetic structure of the studied material; however, its sensitivity is usually limited for quantum dot states. To complement these measurements there is also used photoluminescence excitation spectroscopy, which probes the energetic structure while monitoring directly quantum dot emission.